FlaxSlim™

New Patented Flax Seed Technology

The 3-Way Industry Comparison Blueprint

Performance Metric	Whole / Milled Flaxseed Powder	Standard Flax Extract (e.g., Generic Lignans)	Flaxslim Precision Matrix
Daily Serving Dose	**25g** (Massive, chalky spoonfuls)	500mg – 1,000mg	**2.5g** (Micro-dose)
Primary Mechanism	Passive bulk fiber volume	Basic antioxidant /phytoestrogen support	**Active Multi-Hormone Signaling (L-Cell Trigger)**
GLP-1 Satiety Activation	Weak / Baseline (digestive delay)	Negligible	**+36% Circulating GLP-1** (Multi-pathway surge)
Weight Loss Efficacy	Moderate, high dosages	Negligible	**60% more total weight loss** from 1/10th the material
Body Composition Focus	General weight loss (includes muscle mass loss)	General antioxidant health	**Pure Fat Shifting:** 86% of total weight loss is pure fat mass
Manufacturing Formats	Bulk bags, whole foods, baking mixes	Tablets, hard-shell capsules	**Premium Formats**: Stick packs, daily gummies, clear liquids, capsules
Weight loss observed in clinical studies	~ 1-3 kg	No convincing effect	4.96 kg over 12 weeks (5.8%)

Backed by Dr. Astrup's initial discoveries and eight subsequent human trials, flax mucilage is proven to be the active compound driving appetite suppression, fat reduction, and weight loss.

Whole Flaxseed / Flax Mucilage

Whole flaxseed	Flax mucilage (Active Compound)
Flax mucilage (Active Compound)	Concentrated soluble fiber action
Clinically documented metabolic benefits	More potent satiety & gastrointestinal effects per gram

Clinical Results & Scientific Substantiation

1. Weight Management & Body Composition

- **Primary Action**: Supports long-term weight management and healthy body composition by actively targeting fat mass.
- **Clinical Evidence**: In a 12-week clinical study, participants taking this specific flax mucilage achieved a statistically significant 5.8% reduction in total body weight, resulting in a 200% greater weight-loss effect than the placebo group (4.96 kg lost vs. 1.33 kg). Of the total weight lost, approximately 86% was derived from fat mass (as measured by Bioelectrical Impedance Analysis), preserving metabolically active lean tissue. Additionally, the intervention encouraged a mean reduction of 4.44 cm in waist circumference and 3.53 cm in hip circumference over 12 weeks.

  (Supported by Reference 8)

2. Appetite Regulation & Caloric Intake

- **Primary Action**: Naturally curbs appetite and promotes early satiety to help reduce spontaneous caloric intake.
- **Clinical Evidence**: Shown in an acute clinical trial to produce a statistically significant increase in the sensation of fullness by roughly 30% at 120 minutes post-consumption, driving a 9% reduction in mealtime energy consumption during a subsequent unrestricted meal compared to placebo.
- **Delivery Form Versatility**: Shown to offer lifestyle flexibility, as easy-to-swallow tablets provided a near-identical satiety and caloric-reduction response to liquid fiber drinks in clinical testing (differing by an insignificant 1% to 4%).

  (Supported by Reference 6)

3. Satiety Hormone Activation & Energy Balance

- **Primary Action**: Naturally curbs appetite by safely regulating a broad spectrum of gut satiety signals simultaneously.
- **Clinical Evidence**: In clinical testing, the active viscous fiber successfully suppressed the primary hunger hormone, ghrelin, for a prolonged duration of 5 to 7 hours post-consumption. Furthermore, it promoted post-meal satisfaction by significantly elevating plasma concentrations of the satiety peptide PYY and naturally boosting active GLP-1 by up to 36% versus an isocaloric placebo.

  (Supported by References 3, 7)

4. Glycemic Control & Metabolic Efficiency

- **Primary Action**: Promotes healthy glucose metabolism by structurally slowing carbohydrate digestion. This helps smooth out the normal glycemic curve during the critical 45- to 60-minute window following a meal, minimizing sharp post-meal fluctuations in blood glucose, sparing systemic insulin demands, and reducing hyperinsulinemia.
- **Clinical Evidence**: Shown in acute clinical testing to help maintain optimal cellular insulin sensitivity already within a healthy baseline, demonstrating a clinically significant 20% to 27% reduction in postprandial insulin area under the curve (AUC).

  (Supported by References 3, 7)

5. Lipid Metabolism & Cardiovascular Health

- **Primary Action**: The physical fat-binding matrix helps moderate the absorption of dietary lipids, supporting healthy postprandial circulating triglyceride levels compared to low-fiber diets. In addition, plasma LDL-cholesterol is reduced by 10-15%,
- **Prolonged Lipid Balance**: Extends post-meal metabolic support by helping to regulate the normal circulation of free fatty acids over a prolonged 7-hour digestive period.

  (Supported by Reference 3)

6. Digestive Health & Intestinal Regularity

- **Primary Action**: Clinically shown to optimize healthy bowel movement patterns and significantly improve gastrointestinal regularity markers.
- **Clinical Evidence**: Features a natural fat-binding matrix that was shown to multiply daily fecal fat excretion by roughly 55%. Additionally, it significantly improved gastrointestinal regularity markers by utilizing water-holding capacity to multiply fecal dry mass output by over 35%.

  (Supported by References 1, 5)

7. Gut Microbiome Optimization & SCFA Activation

- **Primary Action**: Acts as a highly fermentable prebiotic substrate, undergoing targeted colonic fermentation to stimulate the natural production of beneficial short-chain fatty acids (SCFAs) like butyrate, which serve as the primary energy source for a healthy intestinal lining and natural gut barrier integrity.
- **Gut-to-Brain Satiety Axis**: The prebiotic fermentation of mucilage into SCFAs helps naturally sustain the extended release of endogenous satiety signals (such as GLP-1 and PYY).
- **Clinical Evidence**: Shown in a 6-week clinical trial to serve as an advanced prebiotic stimulant, driving a pronounced increase in the beneficial Bifidobacterium and the healthy gut-barrier strain Akkermansia, while favorably altering the relative abundance of 33 distinct metagenomic bacterial species.

  (Supported by References 7, 10)

Attached Study References

- **Reference 1**: Kristensen M, et al. (2008). Whole flaxseeds but not sunflower seeds in rye bread reduce apparent digestibility of fat in healthy volunteers. European Journal of Clinical Nutrition, 62(8), 961–967.
- **Reference 2**: Kristensen M, et al. (2013). Linseed Dietary Fibers Reduce Apparent Digestibility of Energy and Fat and Weight Gain in Growing Rats. Nutrients, 5(8), 3287–3298.
- **Reference 3**: Kristensen M, et al. (2012). Flaxseed dietary fibers suppress postprandial lipemia and appetite sensation in young men. Nutrition, Metabolism and Cardiovascular Diseases, 23(2), 136–143.
- **Reference 4 & 5**: Kristensen M, et al. (2012). Flaxseed dietary fibers lower cholesterol and increase fecal fat excretion, but magnitude of effect depend on food type. Nutrition & Metabolism, 9(1), 8.
- **Reference 6**: Ibrügger S, et al. (2012). Flaxseed dietary fiber supplements for suppression of appetite and food intake. Appetite, 58(2), 490–495.
- **Reference 7**: Lafond DW, et al. (2015). Effects of Two Dietary Fibers as Part of Ready-to-Eat Cereal (RTEC) Breakfasts on Perceived Appetite and Gut Hormones in Overweight Women. Nutrients, 7(2), 1245–1266.
- **Reference 8**: Bongartz U, et al. (2022). Flaxseed Mucilage (IQP-LU-104) Reduces Body Weight in Overweight and Moderately Obese Individuals in a 12-week, Three-Arm, Double-Blind, Randomized, and Placebo-Controlled Clinical Study. Obesity Facts, 15(3), 395–404.
- **Reference 10**: Brahe LK, et al. (2015). Dietary modulation of the gut microbiota–a randomised controlled trial in obese postmenopausal women. Nutrition & Diabetes, 5(6), e159

Summary of Material Consistency

Study	Same material/specification as later mucilage studies?	Comment
Study 1	No	Whole flaxseed used; discovery study identifying physiological effect
Study 2	No	Mechanistic in vitro study used to identify active fraction
Study 3	Yes	Purified flaxseed mucilage meeting established compositional specifications
Study 4	Yes	Rat study using purified mucilage matched to established specification
Studies 5 - 6	Yes	Same purified mucilage specification
Study 7 (GLP-1 cereal study)	Yes	Flax fiber extract from same supplier consistent with the established mucilage specification
Studies 8 - 10	Yes	Purified mucilage matched to the same compositional and soluble-fiber specifications

This statement has not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease. "Not all of the claims supported by the scientific studies are permitted under the laws and regulations of all countries. Customers should seek legal counsel for each country in which the customer intends to market and sell these products to assure that the advertising Results made are permitted."

